

Almost upper directed Markov chains on trees


Luis Fredes

(Work with J.F. Marckert)

IMT


Transition matrix: $M = [M_{i,j}]_{i,j \in S}$ with non-negative real entries that sum up to one on each row. A Markov chain Y with transition matrix M satisfies

$$\mathbb{P}(X_{n+1} = b | X_n = a) = M_{a,b}.$$



Transition matrix: $M = [M_{i,j}]_{i,j \in S}$ with non-negative real entries that sum up to one on each row. A Markov chain Y with transition matrix M satisfies

$$\mathbb{P}(X_{n+1} = b | X_n = a) = M_{a,b}.$$



Irreducible: every pair $a, b \in S$ has a finite sequence $\ell = \ell(a, b)$ of steps with positive probability ($M_{a,b}^\ell > 0$) such that b can be reached from a .

From now on we assume irreducibility

Recurrence / Transience: A chain with transition matrix M is called recurrent if for all/one state $a \in S$ the probability to return to a is one, otherwise the chain is called transient.


Positive recurrence: The expected return time of all/one state $a \in S$ is finite, i.e. $\mathbb{E}_a(\tau_a^+) < +\infty$.

Invariant measure: A measure π on S is said to be invariant by M if it is positive and


$$\sum_{a \in S} \pi_a M_{a,b} = \pi_b \quad \text{for all } b \in S.$$

Almost upper and lower directed markov chains on trees


Almost-upper directed \triangleleft and almost-lower directed \triangleright




Nearest neighbors random walks are both



Almost-upper directed \square and almost-lower directed \square on \mathbb{N}



Birth and death processes (BDP) are both



Almost-upper directed \sqsupset and almost-lower directed \sqsubset on \mathbb{N}

$$U := \begin{bmatrix} U_{0,0} & U_{0,1} & U_{0,2} & U_{0,3} & U_{0,4} & \cdots \\ U_{1,0} & U_{1,1} & U_{1,2} & U_{1,3} & U_{1,4} & \cdots \\ 0 & U_{2,1} & U_{2,2} & U_{2,3} & U_{2,4} & \cdots \\ 0 & 0 & U_{3,2} & U_{3,3} & U_{3,4} & \cdots \\ \vdots & \vdots & \vdots & \ddots & \ddots & \cdots \end{bmatrix}, \quad L := \begin{bmatrix} L_{0,0} & L_{0,1} & 0 & 0 & 0 & \cdots \\ L_{1,0} & L_{1,1} & L_{1,2} & 0 & 0 & \cdots \\ L_{2,0} & L_{2,1} & L_{2,2} & L_{2,3} & 0 & \cdots \\ L_{3,0} & L_{3,1} & L_{3,2} & L_{3,3} & L_{3,4} & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots & \cdots \end{bmatrix}$$

Birth and death processes (BDP)

$$T = \begin{bmatrix} T_{0,0} & T_{0,1} & 0 & 0 & 0 & \cdots \\ T_{1,0} & T_{1,1} & T_{1,2} & 0 & 0 & \cdots \\ 0 & T_{2,1} & T_{2,2} & T_{2,3} & 0 & \cdots \\ 0 & 0 & T_{3,2} & T_{3,3} & T_{3,4} & \cdots \\ \vdots & \vdots & \vdots & \ddots & \ddots & \cdots \end{bmatrix}$$

Theorem: Birth and death process (Karlin & McGregor '57)

The following measure π with $\pi_0 = 1$

$$\pi_a = \prod_{j=1}^a \frac{T_{j-1,j}}{T_{j,j-1}} \quad \text{for all } a \geq 1,$$

is the **unique** invariant by T up to a constant factor and the chain is

- **positive recurrent** if and only if

$$\sum_{k \geq 1} \prod_{j=1}^k \frac{T_{j-1,j}}{T_{j,j-1}} < +\infty$$

- **recurrent** if and only if

$$\sum_{k \geq 0} \prod_{j=1}^k \frac{T_{j,j-1}}{T_{j,j+1}} = +\infty.$$

Theorem: Almost-upper triangular case on \mathbb{N} (F. -Marckert '21)

The following measure π with $\pi_0 = 1$

$$\pi_a := \frac{\det(\text{Id} - U_{[0,a-1]})}{\prod_{j=1}^a U_{j,j-1}} \quad \text{for all } a \geq 1.$$

is the **unique** invariant by U up to a constant factor and the chain is

- **positive recurrent** if and only if π_u could be renormalize into a probability distribution.
- **recurrent** if and only if

$$\lim_{b \rightarrow +\infty} U_{1,0} \frac{\det(\text{Id} - U_{[2,b-1]})}{\det(\text{Id} - U_{[1,b-1]})} = 1.$$

Upper directed auxiliary Kernel on a tree




Figure: Construction of the auxiliary transition matrix.

Upper directed auxiliary Kernel on a tree




Figure: Transitions from nodes on $[\emptyset, u]$ are redirected to the common ancestor with u .

Upper directed auxiliary Kernel on a tree




Figure: The total mass associated to the transition.

Upper directed auxiliary Kernel on a tree




Figure: The total mass as a difference of two trees.

Upper directed auxiliary Kernel on a tree

Define :

- The parent of u as $p(u)$.
- The set of nodes belonging to the unique path from \emptyset to u as $[\![\emptyset, u]\!]$.
- For a subset of nodes B , the mass going to B as $U_{a,B} = \sum_{b \in B} U_{a,b}$.
- The tree T_a as the subtree of T rooted at a .
- For a node u in T and $v \in [\![\emptyset, p(u)]\!]$, the successor of v in the direction of u is denoted by $s(v, u)$

Consider the weighted graph $(^uG, ^uU)$ with set of nodes $^uV := [\![\emptyset, u]\!]$, and in which the weight of

$$^uU_{a,b} = (U_{a,T_b} - U_{a,T_{s(b,u)}}) \mathbf{1}_{b \in [\![p(a), p(u)]\!]} + \mathbf{1}_{b=u} U_{a,T_u}$$

is defined for all $a \in [\![\emptyset, p(u)]\!]$ and $b \in [\![\emptyset, u]\!]$

Theorem: Almost upper directed MC on trees (F. -Marckert '24)

The following measure π that we call h -invariant with $\pi_\emptyset = 1$

$$\pi_u := \frac{\det(\text{Id} - {}^u U_{[\emptyset, p(u)]})}{\prod_{v \in [\emptyset, u]} U_{v, p(v)}} \quad \text{for all } u \in T.$$

is **one** invariant measure of U and the chain is

- **positive recurrent** if and only if π could be renormalize into a probability distribution.
- **recurrent** if and only if for any children i of \emptyset

$$\lim_{b \rightarrow +\infty} U_{i, \emptyset} \frac{\det(\text{Id} - U_{[i, < b]}^{(i)})}{\det(\text{Id} - U_{[i, < b]})} = 1.$$

where $U_{[i, < h]}$ is the restriction of U to the subtree T_i to the $h - 1$ first levels of T .

Remark : In the case of recurrence the h -invariant is the unique invariant measure up to constant.

Plan of the talk

- Proof 1/2 : the h -invariant is an invariant measure for U .
- How to compute the h -invariant at a given point.
- The h -invariant as an invariant measure of a conditioned Markov chain.
- Discussion about ends.
- Spectral properties.

Combinatorial warm up I: Matrix tree theorem

$\text{ST}(G)$ = set of spanning trees of G .

Matrix-tree theorem [Kirchhoff]

$$|\text{ST}(G)| = \det \left(\text{Laplacian}_G^{(r)} \right),$$

where $\text{Laplacian}_G^{(r)}$ is the Laplacian matrix of G deprived of the line and column associated to r .

$$\text{Laplacian}_G(i,j) = [\deg(u_i) \mathbb{1}_{i=j} - |\{u_i, u_j\} \in E|]$$

Combinatorial warm up I: Matrix tree theorem


$\text{ST}(G)$ = set of spanning trees of G .

Matrix-tree theorem [Kirchhoff]

$$|\text{ST}(G)| = \det \left(\text{Laplacian}_G^{(r)} \right),$$

where $\text{Laplacian}_G^{(r)}$ is the Laplacian matrix of G deprived of the line and column associated to r .

$$\text{Laplacian}_G(i,j) = [\deg(u_i) \mathbb{1}_{i=j} - |\{u_i, u_j\} \in E|]$$



$$\text{Laplacian}_G = \begin{pmatrix} 3 & -1 & 0 & -1 & 0 & -1 \\ -1 & 5 & -2 & -1 & -1 & 0 \\ 0 & -2 & 3 & 0 & -1 & 0 \\ -1 & -1 & 0 & 3 & -1 & 0 \\ 0 & -1 & -1 & -1 & 4 & -1 \\ -1 & 0 & 0 & 0 & -1 & 2 \end{pmatrix}$$

$$|\text{ST}(G)| = \det \left(\text{Laplacian}_G^{(A)} \right) = 98.$$


Weighted trees

Canonical orientation on rooted trees : all the edges pointing towards the root.

Consequence : every vertex but the root has *exactly one outgoing edge*.

How does a spanning tree of the allowed transitions look like?

Here we work on the graph G whose set of vertices are the same vertices of the tree T and whose set of edges are all having positive transition probability with respect to U .




Weighted trees

Canonical orientation on rooted trees : all the edges pointing towards the root.

Consequence : every vertex but the root has *exactly one outgoing edge*.

How does a spanning tree of the allowed transitions look like?

Here we work on the graph G whose set of vertices are the same vertices of the tree T and whose set of edges are all having positive transition probability with respect to U .



$\text{ST}(G, r)$:= set of spanning trees τ of G (**finite graph**) rooted at r .

Weight of a rooted spanning tree, according to M :

$$W_M(\tau, r) := \prod_{\vec{e} \in \tau} M_{\vec{e}},$$

with edges pointing towards the root r .

Weighted Matrix-tree theorem

$$\sum_{\tau \in \text{ST}(G, r)} W_M(\tau, r) = \det \left(\text{Id} - M^{(r)} \right),$$

where $M^{(r)}$ is the matrix M deprived of the line and column r .


Markov chain tree theorem

The invariant probability measure ρ of M satisfies

$$\rho_v \stackrel{\text{(Cramer)}}{=} \frac{\det(I - M^{(v)})}{Z} \stackrel{\text{(WMTT)}}{=} \frac{\sum_{\tau \in \text{ST}(G, v)} W_M(\tau, v)}{Z}$$

Proof 1/2 : h -invariant for finite trees

We characterize weighted rooted spanning trees τ on the graph of the Markov chain.



A spanning tree τ seen from each vertex gives:

- If a vertex $v \notin [\emptyset, u]$, then it is forced to take the edge $(v, p(v))$. These edges generates a forest.
- Each vertex $v \in [\emptyset, u]$ joins a connected component of the forest and it reaches another vertex in $[\emptyset, u]$.
- We contract the connected components of the forest. This translates the weighted trees of red edges as the determinant of the Laplacian matrix of "U".

Proof 1/2 : h -invariant for finite trees

So the weight of spanning trees rooted at u is given by

$$\sum_{\tau \in \text{ST}(G, r)} W_U(\tau, r) = \det(\text{Id} - {}^u U_{[\emptyset, p(u)]}) \prod_{v \notin [\emptyset, u]} U_{v, p(v)} = C \frac{\det(\text{Id} - {}^u U_{[\emptyset, p(u)]})}{\prod_{v \in [\emptyset, u]} U_{v, p(v)}}$$

From the Markov Chain tree theorem one has that the (unique up to a constant) invariant measure π_u of the chain U at u is proportional to the weight of spanning trees τ rooted at u .

Proof 1/2 : h -invariant for infinite trees


Now for infinite trees, the invariant equations give

$$\rho_u = \sum_{v \in [\emptyset, v]} \rho_v U_{v,u} + \sum_{v \in \text{children}_T(u)} \rho_v U_{v,u} \quad (1)$$

We show that π_u satisfies this system with an auxiliary irreducible Markov chain on a finite tree. The finite tree is the restriction of the tree T until generation $|u| + 1$ and its transition matrix $U|$ coincides with U for transitions up to $|u|$

$$U|_{u,v} = \begin{cases} U_{u,v} & \text{if } |v| < |u| + 1 \\ U_{u,T_v} & \text{if } |v| = |u| + 1 \end{cases}$$

And from the previous part π_u solves the invariance equation (1).



Proof 1/2 : h -invariant for infinite trees


Now for infinite trees, the invariant equations give

$$\rho_u = \sum_{v \in [\emptyset, v]} \rho_v U_{v,u} + \sum_{v \in \text{children}_T(u)} \rho_v U_{v,u} \quad (1)$$

We show that π_u satisfies this system with an auxiliary irreducible Markov chain on a finite tree. The finite tree is the restriction of the tree T until generation $|u| + 1$ and its transition matrix $U|$ coincides with U for transitions up to $|u|$

$$U|_{u,v} = \begin{cases} U_{u,v} & \text{if } |v| < |u| + 1 \\ U_{u,T_v} & \text{if } |v| = |u| + 1 \end{cases}$$

And from the previous part π_u solves the invariance equation (1).



The leaf addition strategy

Let τ and τ' be two infinite or finite trees such that τ' can be obtained from τ by the suppression of a leaf ℓ . Assume that U and U' are two ∇ on τ and τ' satisfying that :

$$U'_{i,j} = \begin{cases} U_{i,j} & \text{if } i \in \tau', j \in \tau' \setminus \{p(\ell)\} \\ U_{i,p(\ell)} + U_{i,\ell} & \text{if } i \in \tau', j = p(\ell) \end{cases}$$

Proposition

If ρ' is a 1-left eigenvector of U' , then ρ given by


$$\rho_i = \begin{cases} \rho'_i & \text{if } i \neq \ell \\ \sum_{v \in [\emptyset, p(\ell)]} \rho'_v \frac{U_{v,\ell}}{U_{\ell,p(\ell)}} & \text{if } i = \ell \end{cases}$$

is a 1-left eigenvector of U .

From our proof we see that " U has the same "h-invariant" information that U on $[\emptyset, u]$

From this we see that we can grow our tree from $\tau' = [\emptyset, p(u)]$ using $U' = {}^p(u)U$ to $\tau = [\emptyset, u]$ with $U = {}^uU$ and they satisfy the leaf addition requirements. This translates into a recursive way to find the value of ρ_u given $\rho_{[\emptyset, p(u)]}$.

$$\rho_u = \sum_{v \in [\emptyset, p(u)]} \rho_v \frac{U_{v, T_u}}{U_{v, p(v)}}$$



Example

Consider a finite tree T and a Markov chain such that starting from u we jump to the parent (when available) or a descendant uniformly at random. We set $\rho_\emptyset = 1$. From the leaf addition applied to $[\![\emptyset, u]\!]$ for $u \neq \emptyset$ we obtain

$$\rho_u = \sum_{v \in [\![\emptyset, p(u)]\!]} \rho_v \frac{U_{v, T_u}}{U_{u, p(u)}} = \sum_{v \in [\![\emptyset, p(u)]\!]} \rho_v \frac{|T_u| / (1_{v \neq \emptyset} + |T_v|)}{1 / (1 + |T_u|)}$$

Now we apply recursively this formula on ρ_v to obtain

$$\rho_u = \frac{|T_u|}{|T_\emptyset|} \left[\prod_{v \in [\![\emptyset, u]\!]} (1 + |T_v|) \right]$$

The h -invariant as the invariant measure of conditioned MC

Let $t \subset T$ a finite subtree of the tree T . Define

$$h_t(u) = \mathbb{P}_u(\tau_t < \infty)$$

Define the Doob's h - transform of the matrix U as the matrix $\tilde{U}(t)$

$$\tilde{U}(t)_{i,j} = \begin{cases} U_{i,j} & \text{if } i \in t \\ \frac{h_t(j)}{h_t(i)} U_{i,j} & \text{if } i \notin t \end{cases}$$

This is the transition matrix of a path conditioned to come back to t .

- The invariant measure π^t of $\tilde{U}(t)$ must coincide with π on the interior of t .
- The invariant measure π^t is unique since $\tilde{U}(t)$ is recurrent.

Conclusion : The measure $\pi^{t_n}(u) \rightarrow \pi(u)$ as $t_n \rightarrow T$ for every $u \in T$.

Meaning that the chain come back to any finite tree

Take home message : the h -measure π of U is the invariant measure of U when “forced to be recurrent”.


When the tree is \mathbb{N} , the system is given by a almost-upper triangular transition matrix, whose balance equations are given by

$$\pi_b = \sum_{a \leq b+1} \pi_a U_{a,b} \iff \pi_{b+1} = (\pi_b - \sum_{a \leq b} \pi_a U_{a,b}) / U_{b+1,b}$$

So in this case the invariant measure is unique (up to a constant factor)

What can we say about ends?

For an infinite path p we define the transition matrix U^p acting on the tree T^p consisting on p together with the finite subtrees hanging from p , such that the jumps of U outside T^p (one ended tree) are considered in U^p as going towards the first common ancestor on T^p .



Proposition

An irreducible \square transition matrix U on an infinite tree T with a finite number of ends is:

- recurrent \iff for every end p , the matrix U^p is recurrent.
- positive recurrent \iff for every end p , the matrix U^p is positive recurrent.

Remark :

- The positive recurrence cares about finite trees, since the chain may spend a lot of time on them.
- To study recurrence, one can prune the finite subtrees and direct all transitions to the first ancestor in an infinite path. The resulting graph is a tree composed with only infinite paths.

What about infinite number of ends?

Lemma

Let U a \square transition matrix on T an infinite tree. If there is an end p with U^p transient, then U is transient.

The reciprocal is not true : we give an example in the complete infinite binary tree, where U^p is positive recurrent for every end p , but the chain U is transient.

Is the number of invariant measures related with the number of ends having a transient U^p ?

...Long story short : The 1-left eigenspace has dimension related with the number of bifurcations in the tree induced by just ends (when dropping finite trees). This bounds the number of extremal invariant measures (only the positive ones are kept).

Is the number of invariant measures related with the number of ends having a transient U^p ?

...Long story short : The 1-left eigenspace has dimension related with the number of bifurcations in the tree induced by just ends (when dropping finite trees). This bounds the number of extremal invariant measures (only the positive ones are kept).

Can we say something about other eigenvalues?

Is the number of invariant measures related with the number of ends having a transient U^p ?

...Long story short : The 1-left eigenspace has dimension related with the number of bifurcations in the tree induced by just ends (when dropping finite trees). This bounds the number of extremal invariant measures (only the positive ones are kept).

Can we say something about other eigenvalues?

Proposition

Let U be a \square irreducible transition matrix on a tree T with no leaves. For all $\lambda \in \mathbb{C}$, the vector $\pi^{(\lambda)}$ defined by $\pi^{(\lambda)}(\emptyset) = 1$ and

$$\pi^{(\lambda)}(u) := \pi^{(\lambda)}(\emptyset) \frac{\det((\lambda \text{Id} - {}^u U)^{(u)})}{\prod_{v \in \llbracket \emptyset, u \rrbracket} U_{v, p(v)}}, \quad \text{for } u \in T$$

is a λ -left eigenvector of U ; hence the spectrum of U is \mathbb{C} .

Thank you for your attention!

Invariant measure and positive recurrence criteria: tridiagonal case

The uniqueness of the invariant measure with $\pi_0 = 1$ gives the equality of two formulas for the invariant measure (Karlin-McGregor and F.-Marckert)

$$\frac{\det(\text{Id} - T_{[0,a-1]})}{\prod_{j=1}^a T_{j,j-1}} = \prod_{j=1}^a \frac{T_{j-1,j}}{T_{j,j-1}} \quad \forall a \geq 1$$

then, the positive recurrence criteria is recovered.

Recurrence criteria

Set $D_{i,j} = \det(\text{Id} - \mathbf{T}_{[i,j]})$, then

$$D_{i,j} = (1 - \mathbf{T}_{i,i})D_{i+1,j} - \mathbf{T}_{i,i+1}\mathbf{T}_{i+1,i}D_{i+2,j}, \quad (2)$$

and set

$$Z_{i,j} := \frac{D_{i,j}}{D_{i+1,j}\mathbf{T}_{i,i-1}}$$

and notice that our result translates into: the MC with transition matrix \mathbf{T} is recurrent iff

$$\lim_{b \rightarrow +\infty} \mathbf{T}_{1,0} \frac{\det(\text{Id} - \mathbf{T}_{[2,b-1]})}{\det(\text{Id} - \mathbf{T}_{[1,b-1]})} = \lim_{b \rightarrow +\infty} \frac{1}{Z_{1,b-1}} = 1.$$

(2) rewrites

$$Z_{i,j} = \frac{(1 - \mathbf{T}_{i,i})}{\mathbf{T}_{i,i-1}} - \frac{\frac{\mathbf{T}_{i,i+1}}{\mathbf{T}_{i,i-1}}}{Z_{i+1,j}},$$

which gives the convergents of a continued fraction.

$$Z_{1,b-1} := c_1 + \cfrac{a_2}{c_2 + \cfrac{a_3}{\ddots + \cfrac{\ddots}{c_{b-2} + \cfrac{a_{b-1}}{c_{b-1}}}}}$$

These can be solved and give that

$$Z_{1,b-1} = 1 + \left(\sum_{k=1}^{b-1} \prod_{j=1}^k \frac{T_{j,j-1}}{T_{j,j+1}} \right)^{-1}$$

So that $\lim_{b \rightarrow \infty} Z_{1,b-1} = 1$ is equivalent to Karlin & McGregor's recurrence criteria.

$$\sum_{k \geq 0} \prod_{j=1}^k \frac{T_{j,j-1}}{T_{j,j+1}} = +\infty.$$